


WO 01/02407

PCT/EP00/06185

28

CLAIMS

1. Compound of formula (I):



2. A process for the preparation of the compound of formula (I), in which 13-(N-Boc-β-isobutylisoserinyl)-14β-hydroxy-baccatine III 1,14-carbonate is either treated with DBU (diazabicyclo[5.4.0] 7-undecene) in methanol or THF, or alternatively is left in solution with methylene chloride or chlorinated solvents in the presence of aliphatic alcohols selected from methanol, ethanol or propanol and with basic alumina, for a time ranging from one hour to 14 days.

3. A process for the preparation of 13-(N-Boc-β-isobutylisoserinyl)-14β-hydroxy-baccatine III or V 1,14-carbonate, which comprises the following steps:

- transformation of 14β-hydroxy-10-deacetylbaccatine III or V into the triethylsilylated derivative at the 7-position;
- preparation of the 1,14 carbonate derivative from the product of step (a);
- selective acetylation of the hydroxyl at 10;
- reaction of the product of step (c) with (4S,5R)-N-Boc-2-(2,4-dimethoxyphenyl)-4-isobutyl-1-oxazolidine-5-

AMENDED SHEET

WO 01/02407

PCT/EP00/06185

29

e) cleavage of the triethylsilyl and dimethoxybenzylidene protective groups from the product of step (d).

4. A process as claimed in claim 3, in which:

the silylating agent of step (a) is triethyl chlorosilane;
the 1,14 carbonate derivative in step (b) is prepared using
phosgene in toluene in methylene chloride/pyridine 3:1
solution under nitrogen atmosphere; the reduction of step
(c) is carried out with LiHMDS in anhydrous THF, and the
resulting 10-hydroxy derivative is subsequently acetylated
with acetyl chloride; the condensation reaction of step (d)
is carried out in anhydrous apolar organic solvent, in the
presence of a base and of the condensing agent
dicyclohexylcarbodiimide (DCC); the triethylsilyl
protective group in step (e) is removed with pyridinium
fluoride in acetonitrile/pyridine solution under nitrogen,
and the dimethoxybenzylidene protective group is removed in
methylene chloride solvent by addition of HCl in methanol
and subsequently of NaHCO_3 .

5. A process for the preparation of 13-(N-Boc- β -isobutylisoserinyl)-14 β -hydroxy-baccatine III or V 1,14-carbonate, which comprises the following steps:

a') selective acetylation of the hydroxyl at C-10 of 14 β -hydroxy-10-deacetyl baccatine III or V;

b') preparation of the 1,14 carbonate derivative from the product of step (a');

c') silylation of the hydroxyl at C-7;

d') reaction of the product of step (c) with (4S,5R)-N-Boc-2-(2,4-dimethoxyphenyl)-4-isobutyl-1-oxazolidine-5-carboxylic acid;

e') cleavage of the triethylsilyl and dimethoxybenzylidene protective groups from the product of step (d').

6. A process as claimed in claim 5, in which the selective acetylation of step (a') is carried out with

ac

scandium,

AMENDED SHEET

WO 01/02407

PCT/EP00/06185

30

ytterbium salts, preferably $\text{CeCl}_3 \cdot 7\text{H}_2\text{O}$, and steps (b')-(e') are carried out analogously to steps (b), (a), (d) and (e) of claim 4.

7. A process for the preparation of (4S, 5R)-N-Boc-2-(2,4-dimethoxyphenyl)-4-isobutyl-1-oxazolidine-5-carboxylic acid, which comprises the following steps:

- a) protection of the amino group of leucinol with Boc;
- b) transformation of N-Boc-L-leucinol into N-Boc-L-leucinal;
- 10 c) preparation of the cyanhydrin of the product of step (b);
- d) transformation of the cyanhydrine nitrile into the corresponding carboxylic acid;
- e) formation of the carboxylic acid methyl ester;
- 15 f) purification of the (2R, 3S)-3-(N-Boc)amino-2-hydroxy-5-methylhexanoic acid methyl ester;
- g) condensation of the product of step (f) with 2,4-dimethoxybenzaldehyde dimethyl acetal;
- h) transformation of (4S, 5R)-N-Boc-2-(2,4-dimethoxyphenyl)-4-isobutyl-1-oxazolidine-5-carboxylic acid methyl ester into the corresponding carboxylic acid.
- 20

8. The following synthesis intermediates: 14 β -hydroxy baccatine III or V, 14 β -hydroxy baccatine III or V 1,14 carbonate, 14- β -hydroxy-7-Tes-10-deacetyl baccatine III or V, 14- β -hydroxy-7-Tes-baccatine ~~III or V~~, 14- β -hydroxy-7-Tes-baccatine III or V 1,14-carbonate, (4S,5R)-N-Boc-2-(2,4-dimethoxyphenyl)-4-isobutyl-1-oxazolidine-5-carboxylic acid.

9. Pharmaceutical compositions containing compound (I) together with pharmaceutically acceptable carriers and excipients.

10. The use of compound (I) for the preparation of a drug with anticancer activity.

AAH

AMENDED SHEET